

THE CENTRAL LIMIT THEOREM FOR LS ESTIMATOR IN SIMPLE LINEAR EV REGRESSION MODELS

Yu Miao ¹, Guangyu Yang ² and Luming Shen ³

¹ Department of Mathematics and Statistics

Wuhan University

Hubei, China 430072

College of Mathematics and Information Science

Henan Normal University

Henan, China 453007

yumiao728@yahoo.com.cn

² Department of Mathematics and Statistics

Wuhan University

Hubei, China 430072

study_yang@yahoo.com.cn

³ Science College

Hunan Agriculture University

Hunan, China 410028

shenluming_20@163.com

Key Words: LS estimator; EV regression models; central limit theorem.

2000 MR Subject Classification: Primary 62F12, Secondary 60F05.

ABSTRACT

In this paper, we obtain the central limit theorems for LS estimator in simple linear errors-in-variables (EV) regression models under some mild conditions. And we also show that those conditions are necessary in some sense.

1. INTRODUCTION

The problem for convergence control of estimator is important in practice. The central limit theorem plays a main role for researching large sample problems. There are numerous studies on the convergence in distribution and various estimation of deviation probability are known, see e.g. Ibragimov and Has'minskii (1979), Ibragimov and Radavicius (1981). Especially, for classical maximum likelihood estimator, Ibragimov and Has'minskii (1979) succeeded in proving the central limit theorem of MLE under some regularity conditions.

In many economic applications, individual observations are very naturally categorized into mutually exclusive and exhaustive groups. For example, individuals can be classified into cohorts and workers are employees of a particular firm. The simplest grouping estimator involves taking the means of all variables for each group and then carrying out a group-level regression by OLS or weighted least squares (if there are different numbers of observations in different groups). This estimator has been called the efficient Wald estimator (cf. Angrist (1991)).

It is well known that there is small sample bias in the efficient Wald estimator because the sample group means do not equal the underlying population means. Deaton (1985) proposed an Errors-in-Variables (EV) estimator to correct the effect of sampling errors.

The main purpose of this paper is to study the asymptotic normality of LS estimator for an EV model. For simplicity of representation, as Liu and Chen (2005), we restrict ourselves to the case of simple linear model:

$$\eta_i = \theta + \beta x_i + \varepsilon_i, \quad \xi_i = x_i + \delta_i, \quad 1 \leq i \leq n \quad (1)$$

with the following assumptions:

- (1) $\theta, \beta, x_1, x_2, \dots$ are unknown constants (parameters);
- (2) $(\varepsilon_1, \delta_1), (\varepsilon_2, \delta_2), \dots$ are independent identically distributed (i.i.d.) random vectors, $\varepsilon_1, \varepsilon_2, \dots$ are i.i.d., $\delta_1, \delta_2, \dots$ are i.i.d., and

$$\mathbb{E}\delta_1 = \mathbb{E}\varepsilon_1 = 0, \quad 0 < Var(\delta_1) = \sigma_1^2, \quad Var(\varepsilon_1) = \sigma_2^2 < \infty;$$

- (3) $\xi_i, \eta_i, i = 1, 2, \dots$ are observable.

From (1) we have

$$\eta_i = \theta + \beta\xi_i + \nu_i, \quad \nu_i = \varepsilon_i - \beta\delta_i, \quad 1 \leq i \leq n. \quad (2)$$

Recently the studies for EV model have attracted much attention: see, for example, Gleser (1981) obtained some large sample results of estimation in a multivariate "errors in variables" regression model. Amemiya and Fuller (1984) discussed the estimation for the multivariate errors-in-variables model with estimated error covariance matrix. Cui (1997) proved the asymptotic normality of M-estimates in the EV model and Liu and Chen (2005) gave the consistency of LS estimator of linear EV regression model under rather weakly conditions and discovered that while in the ordinary linear regression (with the errors i.i.d., the error-variance is finite and non-zero) the weak, strong and quadratic-mean consistency of the regression estimates are equivalent, it is not so in the EV case, i.e., the quadratic-mean consistency requires much more stronger conditions.

Consider formally (2) as a usual regression model of η_i on ξ_i , we get the LS estimator of θ and β as

$$\hat{\beta}_n = \frac{\sum_{i=1}^n (\xi_i - \bar{\xi}_n)(\eta_i - \bar{\eta}_n)}{\sum_{i=1}^n (\xi_i - \bar{\xi}_n)^2}, \quad \hat{\theta}_n = \bar{\eta}_n - \hat{\beta}_n \bar{\xi}_n, \quad (3)$$

where $\bar{\xi}_n = n^{-1} \sum_{i=1}^n \xi_i$, and other similar notations, such as $\bar{\eta}_n$, $\bar{\delta}_n$, are defined in the same way.

Under some common assumptions, Liu and Chen (2005) proved the consistency of LS estimators of simple linear EV model, and obtained that both weak and strong consistency of the estimator are equivalent, but it is not so for quadratic-mean consistency. They also proved that the following condition is the sufficient and necessary condition for $\hat{\beta}_n$ being strong and weak consistent estimate of β :

$$\lim_{n \rightarrow \infty} n^{-1} S_n = \infty,$$

where $S_n = \sum_{i=1}^n (x_i - \bar{x}_n)^2$.

2. THE CENTRAL LIMIT THEOREM FOR THE ESTIMATOR OF PARAMETER β

The model to be studied can be exactly described as follows:

$$(C) : \begin{cases} \eta_i = \theta + \beta x_i + \varepsilon_i, \xi_i = x_i + \delta_i, 1 \leq i \leq n; \\ (\varepsilon_i, \delta_i), 1 \leq i \leq n, i.i.d.; \\ \mathbb{E}\delta_1 = \mathbb{E}\varepsilon_1 = 0, \mathbb{E}\delta_1^2 = \sigma_1^2, \mathbb{E}\varepsilon_1^2 = \sigma_2^2, 0 < \sigma_1^2, \sigma_2^2 < \infty. \end{cases} \quad (4)$$

Here (ξ_i, η_i) , $1 \leq i \leq n$ are observable, while $x_i, 1 \leq i \leq n$, $\theta, \beta, \sigma_1^2, \sigma_2^2$ are unknown parameters. We are mainly interested in the convergence rate of the estimators for the regression coefficient β and constant term θ .

Before our works, we mention a key lemma.

Lemma 1. (*Feller (1971), p530*) For each $n \geq 1$, let $X_{n,1}, \dots, X_{n,n}$ be n independent variables with distribution $F_{n,k}$, $1 \leq k \leq n$. Let $T_n = \sum_{k=1}^n X_{n,k}$. Suppose that $\mathbb{E}(X_{n,k}) = 0$, $\mathbb{E}(T_n^2) = 1$, and that

$$\sum_{k=1}^n \int_{|x|>r} x^2 F_{n,k}(dx) \rightarrow 0 \quad (5)$$

for each $r > 0$. Then the distribution of T_n tends to $N(0,1)$, which is a standard normal distribution.

Theorem 1. Under model (4), let S_n denote $\sum_{i=1}^n (x_i - \bar{x}_n)^2$ and assume that the following condition satisfied:

$$\lim_{n \rightarrow \infty} \frac{n}{\sqrt{S_n}} = 0. \quad (6)$$

In addition, let the following conditions be fulfilled: there exists a constant $\alpha > 0$, such that $\mathbb{E}|\varepsilon_i|^{2+\alpha} < \infty$, $\mathbb{E}|\delta_i|^{2+\alpha} < \infty$ and

$$\lim_{n \rightarrow \infty} \max_{1 \leq i \leq n} \frac{|x_i - \bar{x}_n|}{S_n^{1/2}} = 0, \quad (7)$$

then $\hat{\beta}_n - \beta$ satisfies the asymptotic normality, i.e.,

$$\frac{\sqrt{S_n}}{\sqrt{\text{Var}(\varepsilon_1 - \beta\delta_1)}}(\hat{\beta}_n - \beta) \xrightarrow{d} N(0,1), \quad (8)$$

where $N(0,1)$ is a standard normal distribution.

Remarks 1. Since $S_n = \sum_{i=1}^n (x_i - \bar{x}_n)^2$, and together with (6) and (7), we have

$$n \ll \sqrt{S_n} \ll \max_{1 \leq i \leq n} |x_i - \bar{x}_n|^2 \ll S_n, \quad (9)$$

where " $a_n \ll b_n$ " denotes $a_n \rightarrow \infty$, $b_n \rightarrow \infty$ and $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 0$. This implies that the LS estimator possess good large-sample properties only when $\{x_n, n \geq 1\}$ has greater dispersion. For instance, if $\{x_n, n \geq 1\}$ is a sequence of i.i.d. random variables with common distribution $N(0, 1)$, the LS estimates have no consistency and asymptotic normality. In Section 4, we will show that the condition (6) is necessary in some sense.

Proof. By simple calculation, we have

$$\hat{\beta}_n = \beta + \frac{1}{\sum_{i=1}^n (\xi_i - \bar{\xi}_n)^2} \left(\sum_{i=1}^n (\xi_i - \bar{\xi}_n) \varepsilon_i - \beta \sum_{i=1}^n (x_i - \bar{x}_n) \delta_i - \beta \sum_{i=1}^n (\delta_i - \bar{\delta}_n)^2 \right). \quad (10)$$

For obtaining our result, we need the following lemmas.

Lemma 2. For any $r > 0$, we have

$$\lim_{n \rightarrow \infty} \mathbb{P} \left(\frac{\sum_{i=1}^n (\delta_i - \bar{\delta}_n)^2}{\sqrt{S_n}} \geq r \right) = 0. \quad (11)$$

Proof. For any $r > 0$, because of the following relation,

$$\mathbb{P} \left(\frac{\sum_{i=1}^n \delta_i^2}{\sqrt{S_n}} \geq r \right) \leq \mathbb{P} \left(\frac{\sum_{i=1}^{\lfloor \sqrt{S_n} \rfloor} (\delta_i^2 - \sigma_1^2) - \sum_{i=n+1}^{\lfloor \sqrt{S_n} \rfloor} (\delta_i^2 - \sigma_1^2) + n\sigma_1^2}{\sqrt{S_n}} \geq r \right),$$

where $\lfloor S_n \rfloor$ denote the integer part of S_n and $\sigma_1^2 = \mathbb{E}\delta_1^2$, we have

$$\begin{aligned} & \mathbb{P} \left(\frac{\sum_{i=1}^n (\delta_i - \bar{\delta}_n)^2}{\sqrt{S_n}} \geq r \right) \leq \mathbb{P} \left(\frac{\sum_{i=1}^n \delta_i^2}{\sqrt{S_n}} \geq r \right) \\ & \leq \mathbb{P} \left(\frac{\sum_{i=1}^{\lfloor \sqrt{S_n} \rfloor} (\delta_i^2 - \sigma_1^2)}{\sqrt{S_n}} \geq \frac{r}{2} \right) + \mathbb{P} \left(\frac{-\sum_{i=n+1}^{\lfloor \sqrt{S_n} \rfloor} (\delta_i^2 - \sigma_1^2)}{\sqrt{S_n}} \geq \frac{r}{2} - \frac{n\sigma_1^2}{\sqrt{S_n}} \right) \\ & \triangleq \Delta_1 + \Delta_2. \end{aligned}$$

Next we will control Δ_1 and Δ_2 respectively. By the law of large numbers, Δ_1 is convergent to zero in probability. For enough large n , we have

$$\begin{aligned} \Delta_2 &= \mathbb{P} \left(\frac{-\sum_{i=n+1}^{\lfloor \sqrt{S_n} \rfloor} (\delta_i^2 - \sigma_1^2)}{\sqrt{S_n}} \geq \frac{r}{2} - \frac{n\sigma_1^2}{\sqrt{S_n}} \right) \\ &\leq \mathbb{P} \left(\frac{-\sum_{i=n+1}^{\lfloor \sqrt{S_n} \rfloor} (\delta_i^2 - \sigma_1^2)}{\sqrt{S_n} - n} \geq \frac{r}{4} \right). \end{aligned}$$

Since the same reasons with Δ_1 , Δ_2 is convergent to zero in probability. From the discussion above, we obtain the lemma. $\square$

Lemma 3. *For any $r > 0$, we have*

$$\lim_{n \rightarrow \infty} \mathbb{P} \left(\left| \frac{\sum_{i=1}^n (\xi_i - \bar{\xi}_n)^2}{S_n} - 1 \right| \geq r \right) = 0. \quad (12)$$

Proof. In fact, we only need to prove the form $\mathbb{P} \left(\frac{\sum_{i=1}^n (\xi_i - \bar{\xi}_n)^2}{S_n} - 1 \geq r \right)$, and the proof of the other form $\mathbb{P} \left(\frac{\sum_{i=1}^n (\xi_i - \bar{\xi}_n)^2}{S_n} - 1 \leq -r \right)$ is similar. By simple calculation, we have

$$\sum_{i=1}^n (\xi_i - \bar{\xi}_n)^2 = \sum_{i=1}^n (x_i - \bar{x}_n)^2 + 2 \sum_{i=1}^n (x_i - \bar{x}_n)(\delta_i - \bar{\delta}_n) + \sum_{i=1}^n (\delta_i - \bar{\delta}_n)^2,$$

and

$$\sum_{i=1}^n (\xi_i - \bar{\xi}_n)^2 - S_n \leq \frac{r}{2} S_n + \frac{2+r}{r} \sum_{i=1}^n (\delta_i - \bar{\delta}_n)^2.$$

Thereby, we get

$$\mathbb{P} \left(\frac{\sum_{i=1}^n (\xi_i - \bar{\xi}_n)^2}{S_n} - 1 \geq r \right) \leq \mathbb{P} \left(\frac{(2+r) \sum_{i=1}^n (\delta_i - \bar{\delta}_n)^2}{r S_n} \geq \frac{r}{2} \right).$$

Hence, Lemma 3 turns to be obvious due to Lemma 2 $\square$

Lemma 4. *For any $r > 0$, we have*

$$\lim_{n \rightarrow \infty} \mathbb{P} \left(\left| \frac{\sum_{i=1}^n (\delta_i - \bar{\delta}_n) \varepsilon_i}{\sqrt{S_n}} \right| \geq r \right) = 0.$$

Proof. For any $r > 0$, we have

$$\mathbb{P} \left(\left| \frac{\sum_{i=1}^n (\delta_i - \bar{\delta}_n) \varepsilon_i}{\sqrt{S_n}} \right| \geq r \right) \leq \mathbb{P} \left(\frac{\sum_{i=1}^n (\delta_i - \bar{\delta}_n)^2}{\sqrt{S_n}} + \frac{\sum_{i=1}^n (\varepsilon_i - \bar{\varepsilon}_n)^2}{\sqrt{S_n}} \geq 2r \right).$$

So, by Lemma 2, our result is obtained. $\square$

Lemma 5. *We have*

$$\frac{\sqrt{S_n}}{\sqrt{\text{Var}(\varepsilon_1 - \beta \delta_1)}} \frac{\sum_{i=1}^n (x_i - \bar{x}_n)(\varepsilon_i - \beta \delta_i)}{S_n} \xrightarrow{d} N(0, 1).$$

Proof. For any $n \geq 1$ and $1 \leq i \leq n$, denote

$$X_{n,i} := \frac{(x_i - \bar{x}_n)}{\sqrt{S_n \text{Var}(\varepsilon_1 - \beta \delta_1)}} (\varepsilon_i - \beta \delta_i).$$

Then for any $n \geq 1$, $\{X_{n,i}\}_{i=1}^n$ is a sequence of independent random variables, and $\mathbb{E}X_{n,i} = 0$, $\sum_{i=1}^n \mathbb{E}X_{n,i}^2 = 1$. Next, we shall consider the following control: for any $r > 0$,

$$\begin{aligned} & \sum_{i=1}^n \mathbb{E}[|X_{n,i}|^2 \mathbf{1}_{\{|X_{n,i}| > r\}}] \\ & \leq \frac{1}{S_n \text{Var}(\varepsilon_1 - \beta \delta_1)} \sum_{i=1}^n \int_{\left\{ \frac{|(x_i - \bar{x}_n)x|}{\sqrt{S_n \text{Var}(\varepsilon_1 - \beta \delta_1)}} > r \right\}} \frac{|(x_i - \bar{x}_n)x|^{2+\alpha}}{r^\alpha [S_n \text{Var}(\varepsilon_1 - \beta \delta_1)]^{\alpha/2}} dF(x), \end{aligned}$$

where $F(x)$ denotes the distribution of $\varepsilon_1 - \beta \delta_1$. Therefore,

$$\begin{aligned} & \sum_{i=1}^n \mathbb{E}[|X_{n,i}|^2 \mathbf{1}_{\{|X_{n,i}| > r\}}] \\ & \leq \frac{\mathbb{E}|\varepsilon_1 - \beta \delta_1|^{2+\alpha}}{r^\alpha (S_n \text{Var}(\varepsilon_1 - \beta \delta_1))^{1+\alpha/2}} \sum_{i=1}^n |x_i - \bar{x}_n|^{2+\alpha} \\ & \leq \frac{\mathbb{E}|\varepsilon_1 - \beta \delta_1|^{2+\alpha}}{r^\alpha (\text{Var}(\varepsilon_1 - \beta \delta_1))^{1+\alpha/2}} \max_{1 \leq i \leq n} \frac{|x_i - \bar{x}_n|^\alpha}{S_n^{\alpha/2}}, \end{aligned}$$

and by (7), we obtain

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \mathbb{E}[|X_{n,i}|^2 \mathbf{1}_{\{|X_{n,i}| > r\}}] = 0.$$

From Lemma 1, we obtain our lemma. □

Now we give the proof of Theorem 1. By (10), we have

$$\sqrt{S_n}(\hat{\beta}_n - \beta) = \sqrt{S_n} \frac{\sum_{i=1}^n (\delta_i - \bar{\delta}_n) \varepsilon_i + \sum_{i=1}^n (x_i - \bar{x}_n) (\varepsilon_i - \beta \delta_i) - \beta \sum_{i=1}^n (\delta_i - \bar{\delta}_n)^2}{\sum_{i=1}^n (\xi_i - \bar{\xi}_n)^2}. \quad (13)$$

From Lemma 3 and Lemma 4, it is easy to see that in probability,

$$\frac{S_n}{\sum_{i=1}^n (\xi_i - \bar{\xi}_n)^2} \frac{\sum_{i=1}^n (\delta_i - \bar{\delta}_n) \varepsilon_i}{\sqrt{S_n}} \rightarrow 0,$$

and with the same reasons, in probability

$$\frac{S_n}{\sum_{i=1}^n (\xi_i - \bar{\xi}_n)^2} \frac{\sum_{i=1}^n (\delta_i - \bar{\delta}_n)^2}{\sqrt{S_n}} \rightarrow 0.$$

Because of Lemma 3 and Lemma 5, we have

$$\frac{S_n}{\sum_{i=1}^n (\xi_i - \bar{\xi}_n)^2} \frac{\sum_{i=1}^n (x_i - \bar{x}_n)(\varepsilon_i - \beta\delta_i)}{\sqrt{S_n}} \xrightarrow{d} N(0, \text{Var}(\varepsilon_1 - \beta\delta_1)).$$

Therefore, the proof of Theorem 1 is completed. $\square$

3. THE CENTRAL LIMIT THEOREM FOR THE ESTIMATOR OF PARAMETER θ

In this section, we shall discuss the central limit theorem of estimator of parameter θ . LS estimate of θ is

$$\hat{\theta}_n = \bar{\eta}_n - \hat{\beta}_n \bar{\xi}_n. \quad (14)$$

Hence

$$\hat{\theta}_n - \theta = (\beta - \hat{\beta}_n) \bar{x}_n + (\beta - \hat{\beta}_n) \bar{\delta}_n - \beta \bar{\delta}_n + \bar{\varepsilon}_n. \quad (15)$$

Liu and Chen (2005), have proved the following theorem, which gave the sufficient and necessary condition for $\hat{\theta}_n$ being weak consistent estimate of θ .

Theorem 2. *Under model (4), a sufficient and necessary condition for the weak consistency of $\hat{\theta}_n$ is*

$$\lim_{n \rightarrow \infty} n \bar{x}_n (S_n^*)^{-1} = 0, \quad S_n^* = \max(n, S_n). \quad (16)$$

Theorem 3. *Under model (4), assume that the conditions of the Theorem 1 are satisfied. In addition, we assume*

$$\frac{S_n}{n \bar{x}_n^2} \rightarrow \infty, \quad (17)$$

then $\hat{\theta}_n - \theta$ satisfies the asymptotic normality, i.e.,

$$\frac{\sqrt{n}}{\sqrt{\text{Var}(\varepsilon_1 - \beta\delta_1)}} (\hat{\theta}_n - \theta) \xrightarrow{d} N(0, 1). \quad (18)$$

Remarks 2. Since $n \bar{x}_n^2 \ll S_n$ and the definition of S_n , we have $\lim_{n \rightarrow \infty} \frac{\sum_{i=1}^n x_i^2}{S_n} = 1$ and (16) satisfied. Furthermore, we notice that the normalization of β_n is related to $\sqrt{S_n}$, however, the normalization of θ is related to $\sqrt{n}$. It is not mysterious, since the parameter β has coefficients x_i , but θ not in EV models. And this also suggests that the proof of Theorem 3 might be easier than that of Theorem 1.

Proof. It is easy to see,

$$\frac{\sqrt{n}}{\sqrt{\text{Var}(\varepsilon_1 - \beta\delta_1)}}(\bar{\varepsilon}_n - \beta\bar{\delta}_n) \xrightarrow{d} N(0, 1).$$

Hence, it is enough to show in probability

$$\frac{\sqrt{n}}{\sqrt{\text{Var}(\varepsilon_1 - \beta\delta_1)}}(\beta - \hat{\beta}_n)(\bar{x}_n + \bar{\delta}_n) \rightarrow 0.$$

By Theorem 1, we only need to show in probability,

$$\frac{\sqrt{n}}{\sqrt{S_n}}(\bar{x}_n + \bar{\delta}_n) \rightarrow 0.$$

The law of large numbers, and together with the condition (17), yields our results. $\square$

4. FURTHER DISCUSSIONS

Recalling that Liu and Chen (2005) proved that the following condition is the sufficient and necessary condition for $\hat{\beta}_n$ being strong and weak consistent estimate of β :

$$\lim_{n \rightarrow \infty} n^{-1}S_n = \infty.$$

However, in Theorem 1, we assume the condition (6), i.e.,

$$\lim_{n \rightarrow \infty} \frac{n}{\sqrt{S_n}} = 0. \tag{19}$$

It seems that the condition (19) is strong. In this section, we will analyze it and show that it is necessary in some sense.

From the proof of Lemma 2, we know that the condition (19) is a technical condition to prove the following limit

$$\lim_{n \rightarrow \infty} \mathbb{P} \left(\frac{1}{\sqrt{S_n}} \sum_{k=1}^n \delta_k^2 \geq r \right) = 0. \tag{20}$$

Before our discussion, we need mention the following result.

Theorem 4. (Petrov, 1987) *Let $\{X_k\}_{k \geq 1}$ is a sequence of independent random variables with the distribution $V_k(x)$ and $\{a_n\}_{n \geq 1}$ is a sequence of increasing positive real number. Then*

$$\frac{1}{a_n} \sum_{k=1}^n X_k \xrightarrow{\mathbb{P}} 0,$$

if and only if the following conditions are satisfied

(i)

$$\sum_{k=1}^n \int_{|x| \geq a_n} dV_k(x) \rightarrow 0;$$

(ii)

$$\frac{1}{a_n^2} \sum_{k=1}^n \left\{ \int_{|x| < a_n} x^2 dV_k(x) - \left(\int_{|x| < a_n} x dV_k(x) \right)^2 \right\} \rightarrow 0;$$

(iii)

$$\frac{1}{a_n} \sum_{k=1}^n \int_{|x| < a_n} x dV_k(x) \rightarrow 0.$$

Under our model, we can take $a_n = \sqrt{S_n}$, $X_k = \delta_k^2$, $k \geq 1$ in Theorem 4. If

$$\frac{1}{\sqrt{S_n}} \sum_{k=1}^n \delta_k^2 \xrightarrow{\mathbb{P}} 0,$$

then, by (iii) of Theorem 4, we have

$$\frac{1}{\sqrt{S_n}} \sum_{k=1}^n \mathbb{E} \delta_k^2 I_{\{\delta_k^2 < \sqrt{S_n}\}} = \frac{n}{\sqrt{S_n}} \mathbb{E} \delta_1^2 I_{\{\delta_1^2 < \sqrt{S_n}\}} \rightarrow 0.$$

Since $0 < \mathbb{E} \delta_1^2 < \infty$, we get

$$\frac{n}{\sqrt{S_n}} \rightarrow 0.$$

That is to say, (6) is the necessary condition for Theorem 1.

5. ACKNOWLEDGEMENTS

The authors wish to thank Prof. L.M. Wu of Université Blaise Pascal and Wuhan University, and Prof. D.H. Hu of Wuhan University for their helpful discussions and suggestions during writing this paper. The second author was partially supported by the National Natural Science Foundation of China (Grant No: 10371092) and the Foundation of Wuhan University. At last, the authors are very grateful to the conscientious anonymous referee for his very serious and valuable report. His suggestions and comments have largely contributed to the Section 4.

BIBLIOGRAPHY

- Akahira, M. and Takeuchi, K. (1981). *Asymptotic efficiency of statistical estimators*. Springer-Verlag New York.
- Amemiya, Y. and Fuller, W.A. (1984). Estimation for the multivariate errors-in-variables model with estimated error covariance matrix. *Ann. Statist.*, **12(2)**, 497-509.
- Angrist, J.D. (1991). Grouped Data Estimation and Testing in Simple Labor Supply Models. *J. Econometrics*, **47**, 243-265.
- Cui, H.J. (1997). Asymptotic normality of M-estimates in the EV model. *J. Sys. Sci. and Math. Sci.*, **10(3)**, 225-236.
- Deaton, A. (1985). Panel data from a time series of cross-sections. *J. Econometrics*, **30**, 109- 126.
- Feller, W. (1971). *An introduction to probability theory and its applications. Second Edition*. John Wiley and Sons, Inc. Vol 2.
- Fuller, W. (1987). *Measurement Error Models*. Wiley, New York.
- Gleser, L. J. (1981). Estimation in a multivariate "errors in variables" regression model: Large sample results. *Ann. Statist.*, **9(1)**, 24-44.
- Ibragimov, I.A. and Has'minskii, R.Z. (1979). *Statistical Estimation*. Springer-Verlag New York.
- Ibragimov, I. and Radavicius, M. (1981). Probability of large deviations for the maximum likelihood estimator. *Soviet Math. Dokl.*, **23(2)**, 403-406.
- Liu, J.X. and Chen, X.R. (2005). Consistency of LS estimator in simple linear EV regression models. *Math. Acta. Sci.*, **25B(1)**, 50-58.
- Petrov, V.V. (1987). *Limit Theorems for Sums of Independent Random Variables*, Nauka, Moscow. (in Russian). English translation, (1991). Oxford Univ. Press, Oxford. Chinese translation, (1991). by Huang, K.M. and Su, C. Technology Press of China, Hefei, China.

Stuart, A., Ord, J. K. and Arnold, S. (1999). *Kendall's Advanced Theory of Statistics*.
Vol.2A. Arnold, London.